

Retinoic Acid Synthesis and Signaling during Early Organogenesis

Gregg Duester^{1,*}

¹Burnham Institute for Medical Research, Development and Aging Program, 10901 North Torrey Pines Road, La Jolla, CA 92037, USA

*Correspondence: duester@burnham.org

DOI 10.1016/j.cell.2008.09.002

Retinoic acid, a derivative of vitamin A, is an essential component of cell-cell signaling during vertebrate organogenesis. In early development, retinoic acid organizes the trunk by providing an instructive signal for posterior neuroectoderm and foregut endoderm and a permissive signal for trunk mesoderm differentiation. At later stages, retinoic acid contributes to the development of the eye and other organs. Recent studies suggest that retinoic acid may act primarily in a paracrine manner and provide insight into the cell-cell signaling networks that control differentiation of pluripotent cells.

Introduction

Retinoic acid (RA) is a small lipophilic molecule derived from vitamin A that stands apart from other diffusible cell-cell signaling factors that direct developmental processes for many reasons. Protein factors, such as fibroblast growth factor (FGF), WNT, hedgehog, or the transforming growth factor beta (TGF- β) superfamily, bind to cell-surface receptors and initiate intracellular signaling pathways. In stark contrast, RA enters the nucleus and binds directly to target genes via nuclear receptors. RA signaling also differs from the major classes of protein growth factors in that it appears to be a chordate invention (Marlétaz et al., 2006). The most convincing evidence that RA signaling is limited to chordates is the observation that only chordates appear to possess the retinaldehyde dehydrogenase (RALDH) enzymes needed to synthesize RA. Thus, during chordate evolution, RA signaling was layered on top of many other pre-existing cell-cell signaling pathways. RA regulates many of the same developmental processes that are controlled by protein growth factors including neurogenesis, cardiogenesis, body axis extension, and development of the forelimb buds, foregut, and eye. In addition, recent studies indicate that RA signaling represses several of these growth factor signaling pathways.

Retinoic Acid Synthesis, Degradation, and Signaling

The ability of vitamin A to influence development is made possible by enzymes controlling the conversion of the alcohol form of vitamin A (retinol) first to an aldehyde (retinaldehyde) and then to a carboxylic acid (retinoic acid; RA) (Figure 1). The first step of RA synthesis, oxidation of retinol to retinaldehyde, is catalyzed by several alcohol dehydrogenases (ADHs) and retinol dehydrogenases (RDHs). Genetic studies suggest that at least three ADHs (ADH1, ADH3, and ADH4) and two RDHs (RDH1 and RDH10) play a physiological role in RA synthesis (Table 1). Expression of these retinol-oxidizing enzymes is widespread and overlapping (Ang et al., 1996; Zhang et al., 2001; Sandell et al., 2007). The second step of RA synthesis, oxidation of retinaldehyde to RA, is catalyzed by three retinaldehyde dehydrogenases (RALDH1, RALDH2, and RALDH3), which display




Figure 1. Retinoic Acid Synthesis and Signaling

Depicted is the paracrine mechanism of retinoic acid (RA) signaling. Retinol is carried in the serum by retinol-binding protein (RBP4) secreted from the liver. Retinol enters cells via a specific receptor STRA6, and cellular retinol-binding protein (CRBP) facilitates conversion of retinol to retinyl esters for storage. In an RA-generating tissue, retinol is oxidized to retinaldehyde by either alcohol dehydrogenase (ADH) or retinol dehydrogenase (RDH), and retinaldehyde is oxidized to RA by retinaldehyde dehydrogenase (RALDH). RA is then released and taken up by surrounding cells. Cells that express cytochrome P450 (CYP26) initiate the further oxidation of RA for degradation and excretion and are not RA target cells. Some RA target cells express cellular RA-binding protein (CRABP) that facilitates uptake of RA and transport to the nucleus where RA binds the RA receptor (RAR). The ternary complex of ligand-bound RAR with RXR and a retinoic acid response element (RARE) regulates transcription of RA target genes by altering the binding of corepressors and coactivators.

Table 1. Enzymes Involved in RA Synthesis during Early Organogenesis in Mouse

Gene	Major Sites of Expression	Knockout Phenotype
Retinaldehyde dehydrogenases (oxidation of retinaldehyde to RA)		
<i>Raldh1</i>	dorsal retina at E9.5	nonlethal; adults fertile; perioptic mesenchyme defect when <i>Raldh3</i> also null; protects against obesity in adult
<i>Raldh2</i>	paraxial mesoderm at E7.5; somites; lateral plate mesoderm; optic vesicle mesenchyme; mesonephros; ventral spinal cord	lethal at E9.5; hindbrain and spinal cord defects; small somites with left-right asymmetry; forelimb bud absent; abnormal heart with medial distended cavity; lung and pancreas agenesis; ventral retina defect when <i>Raldh3</i> also null
<i>Raldh3</i>	lens and nasal placodes at E8.75; retinal pigment epithelium; ventral retina; mesonephros; ventral forebrain	lethal at birth (blockage of nasal passages); ventral retina defect; forebrain striatal defect; perioptic mesenchyme defect when <i>Raldh1</i> also null
Alcohol/Retinol dehydrogenases (oxidation of retinol to retinaldehyde)		
<i>Adh1</i>	mesonephros at E9.5; liver; kidney	nonlethal; adults fertile; hypersensitive to vitamin A toxicity; increased retinyl esters
<i>Adh3</i>	ubiquitous from E6.5 to adult	postnatal growth deficiency; adults fertile but smaller litters; hypersensitive to vitamin A deficiency
<i>Adh4</i>	posterior mesoderm at E7.5; craniofacial mesenchyme; stomach	nonlethal; adults fertile; hypersensitive to vitamin A deficiency
<i>Rdh1</i>	widespread at E7.5; liver; kidney	nonlethal; adults fertile; increased weight and adiposity; increased retinyl esters
<i>Rdh10</i>	lateral plate mesoderm at E7.5; somites; floor plate of neural tube	lethal at E13.0; small forelimbs; craniofacial defects; lung agenesis

nonoverlapping tissue-specific patterns of expression during embryogenesis (Mic et al., 2002) (Table 1). Oxidation of RA, which leads to its degradation, is carried out by three cytochrome P450 (CYP) enzymes known as CYP26A1 (Abu-Abed et al., 2001), CYP26B1 (Yashiro et al., 2004), and CYP26C1 (Uehara et al., 2007). These enzymes also display unique tissue-specific patterns of expression during mouse embryogenesis, suggesting that they influence where RA signaling is able to occur in the embryo.

RA serves as a ligand for two families of nuclear receptors that bind DNA and directly regulate transcription: (1) the RA receptors (RAR α , RAR β , and RAR γ) that bind the abundant form of RA known as all-*trans*-RA and (2) the retinoid X receptors (RXR α , RXR β , and RXR γ) that bind an isomer known as 9-*cis*-RA (Chawla et al., 2001). However, 9-*cis*-RA is normally undetectable except when vitamin A is present in excess (Arnhold et al., 1996; Mic et al., 2003). Hence, it may play a pharmacological but not a physiological role as an RXR ligand. As RXR forms heterodimers with RAR and several other nuclear receptors when bound to DNA, this suggests that RXR functions as a scaffold protein to facilitate DNA binding for several different types of nuclear receptors (Chawla et al., 2001). In vivo studies have demonstrated that ligand binding to just the RAR portion of RAR/RXR heterodimers is sufficient and necessary to rescue a lethal defect in RA synthesis, whereas ligand binding to RXR does not rescue the defect and is unnecessary (Mic et al., 2003). When RA binds to the RAR partner of RAR/RXR heterodimers bound to a regulatory DNA element, this stimulates a cascade of events resulting in recruitment of transcriptional coactivators and initiation of transcription (Germain et al., 2002).

RA is not produced by all cells of the body at all stages of development but is instead produced in a unique spatiotemporal pattern. Retinol is secreted by the liver and transported in the blood at micromolar levels via serum retinol-binding protein (RBP4) and is made available to all cells (including

embryonic cells by maternal transfer) for potential conversion to RA (Quadro et al., 1999). Many cells possess STRA6, which functions as a membrane receptor for RBP4 to facilitate retinol uptake (Kawaguchi et al., 2007). Many cells also contain cellular retinol-binding proteins (CRBPs) that bind retinol inside the cell (Noy, 2000). CRBP1 has been proposed to facilitate conversion of retinol to retinyl esters for storage and to facilitate oxidation of retinol to retinaldehyde by RDHs, but not ADHs, for RA synthesis. These findings suggested that RDHs but not ADHs are important in endogenous RA synthesis. However, recent comparisons of ADH and RDH retinol enzymatic activity in the presence or absence of CRBP1 concluded that free retinol but not CRBP1-bound retinol is the substrate for both enzyme families (Gallego et al., 2006). Also, mice lacking CRBP1 do not exhibit decreased RA synthesis but do have greatly reduced stores of liver retinyl esters and are thus very sensitive to vitamin A deficiency (Ghyselinck et al., 1999). Interestingly, mice lacking ADH1 have higher than normal levels of liver retinyl esters, and mice deficient in both ADH1 and CRBP1 have relatively normal levels of liver retinyl esters and reduced sensitivity to vitamin A deficiency (Molotkov et al., 2004). Thus, ADH1 and CRBP1 have opposing roles in the liver that prevent toxic accumulation of retinol but still enable a large fraction of retinol to become esterified as a stored form, thus providing a continuous source of retinol to be secreted into the blood for use in peripheral tissues including embryos.

Studies on mice carrying null mutations in ADH and RDH enzymes suggest that oxidation of retinol to retinaldehyde may be controlled in vivo by multiple genes (Table 1). Mice lacking ADH1 are susceptible to retinol toxicity yet are otherwise phenotypically normal, indicating that oxidation of retinol by ADH1 contributes to removal of excess retinol (via further metabolism to RA and oxidized forms of RA) rather than contributing to RA needed for signaling (Molotkov et al., 2002a). In contrast, loss of ADH3 and ADH4 impairs postnatal survival during vitamin A deficiency, suggesting potential roles in RA synthesis designed

for RA signaling (Molotkov et al., 2002b). However, given that ADH3 has very low activity for retinol oxidation (Molotkov et al., 2002b), the effect of its knockout may be unrelated to RA synthesis. In comparison, ADH4 and RDH1 are very efficient in retinol oxidation (Zhang et al., 2001; Gallego et al., 2006), yet mice lacking ADH4 and RDH1 have not been shown to display detectable alterations in embryonic RA signaling (Molotkov et al., 2002b; Zhang et al., 2007). Among these genes, only mutants in *Rdh10* have a serious defect in embryonic RA signaling resulting in embryonic lethality at embryonic day 13 (E13) (Sandell et al., 2007). *Rdh10* mutant embryos still maintain RA signaling in some embryonic tissues, suggesting that ADH1, ADH3, ADH4, or RDH1, which are all expressed in embryos, may also function to generate retinaldehyde for RA synthesis.

Retinol oxidation is a reversible reaction. Hence, the ability to convert retinaldehyde back to retinol, a reaction that multiple enzymes can accomplish, may provide further control over RA synthesis (Gallego et al., 2006). In contrast, oxidation of retinaldehyde to RA is irreversible. Given the widespread access to retinol via the circulatory system, it is possible that all cells establish an equilibrium between retinol and retinaldehyde, but only cells expressing one of the RALDHs can oxidize the available retinaldehyde to RA.

Conversion of retinol to RA occurs at relatively low levels, but RA has been detected in mouse embryos using sensitive RA-reporter assays. Evidence from a mouse strain that bears a transgene expressing *lacZ* under the control of a retinoic acid response element (RARE) indicates that RA signaling activity is first observed at E7.5, the late primitive streak stage, and is localized to the trunk from E7.5–E8.5 (Rossant et al., 1991). An RA-reporter cell line was used to demonstrate that mouse embryo explants from E7.5, but not E6.5, have detectable RA activity (Ang et al., 1996). The RA biosynthetic enzyme RALDH2 is first expressed at E7.5 in trunk paraxial mesoderm and by E8.5 displays expression in paraxial and lateral plate mesoderm that appears quite similar to the pattern of RA localization (Sirbu et al., 2005; Molotkova et al., 2005). Studies on mice lacking RALDH2 have shown that it is responsible for all RA signaling activity in the embryo from E7.5 (Sirbu et al., 2005) to E8.5 (Molotkova et al., 2005), but that immediately after this stage RALDH1 and RALDH3 also contribute to RA synthesis in the eye and olfactory pit (Molotkov et al., 2006).

RA released by RA-generating cells can enter adjacent cells where it has two main fates (Figure 1). In cells expressing one of the *Cyp26* genes, RA is degraded and RA signaling is prevented (Hernandez et al., 2007; Uehara et al., 2007). In cells not expressing *Cyp26*, RA can enter the nucleus and bind to RAR receptors leading to transcription of target genes. Additionally, some cells express cellular RA-binding protein-2 (CRABP2), which greatly facilitates cellular uptake of RA and transfer to the nucleus (Sessler and Noy, 2005; Schug et al., 2007).

Retinoic Acid Influences Induction and Patterning of Embryonic Tissues

Vitamin A is unique among the vitamins in that its concentration must be within a very narrow range in order to avoid both deficiency and toxicity. Thus, adding vitamin A or RA to embryos can easily induce teratogenic effects including major alterations

Table 2. Key Target Genes Regulated by Retinoic Acid during Early Organogenesis

Developmental Process	Key Target Genes
Hindbrain anteroposterior patterning	↑ <i>Hoxa1</i> , <i>Hoxb1</i> , <i>Hoxa3</i> , <i>Hoxd4</i> , <i>vHnf1</i>
Spinal cord motor neuron differentiation	↑ <i>Pax6</i> , <i>Olig2</i>
Formation of early somites	↓ <i>Fgf8</i> , ↑ <i>Cdx1</i>
Heart anteroposterior patterning	↓ <i>Fgf8</i>
Forelimb induction	↓ <i>Fgf8?</i>
Pancreas induction	↑ <i>Pdx1</i>
Lung induction	↑ <i>Hoxa5</i> , ↓ <i>TGF-β1</i>
Optic cup formation	?
Anterior eye formation	↑ <i>Pitx2</i>
Kidney formation	↑ <i>Ret</i>
Induction of meiosis	↑ <i>Stra8</i>

Expression (↑) induced or (↓) repressed.

in organogenesis. However studying the teratogenic effects of RA is often not useful for determining its physiological role given that high levels of RA may induce or repress genes not normally regulated by endogenous RA. Thus, loss-of-function studies are necessary to determine the normal function of RA during organogenesis.

Early studies on the embryonic effects of a loss of RA were performed using vitamin A deficiency (Clagett-Dame and DeLuca, 2002; Dersch and Zile, 1993; Dickman et al., 1997) or compound RAR mutations (Lohnes et al., 1994; Mendelsohn et al., 1994). These early studies indicated that RA was essential for development of several organs including the hindbrain, spinal cord, heart, eye, skeleton, forelimb buds, lung, pancreas, and genitourinary tract. More recently, the use of RALDH mutations that completely eliminate RA synthesis in specific tissues at early stages of development has made it possible to examine the mechanism of RA action in detail as described below. Cumulatively, these studies show that RA provides an instructive signal for posterior neuroectoderm (hindbrain, spinal cord) and posterior foregut endoderm (pancreas, lung) and a permissive signal for trunk mesoderm (somites, heart, forelimb) in early development. At later stages RA contributes to the development of the eye and other organs. Defects in the forelimb bud, lung, and pancreas can be characterized as defects in induction of these tissues, as a loss of RA inhibits organogenesis. In contrast, defects in hindbrain, spinal cord, heart, somites, and eye occur after organogenesis has been induced and can be characterized as defects in patterning or morphogenesis of these tissues. Many important RA target genes in these tissues have been found (Table 2).

Retinoic Acid Organizes the Trunk

RA generated by RALDH2 in the presomitic mesoderm is released and travels to the adjacent posterior neuroectoderm and posterior foregut endoderm where it induces various homeobox genes in posterior hindbrain and trunk tissues. This instructive mechanism of RA action is widely accepted as the main role of RA signaling for early organogenesis. However, recent findings indicate that RA can also influence




Figure 2. Retinoic Acid Organizes the Trunk by Repressing *Fgf8*

Shown on the left is an E8.0 mouse embryo stained for *fibroblast growth factor 8* (*Fgf8*) mRNA by in situ hybridization. At this stage *Fgf8* mRNA is expressed in an anterior cardiac domain where it induces *Isl1*, plus a posterior domain encompassing the epiblast and primitive streak (ps) where FGF8 is needed during gastrulation and neural induction; *retinaldehyde dehydrogenase 2* (*Raldh2*) is expressed in between these two domains in the somitic mesoderm where the future trunk will form. The diagram depicts the function of retinoic acid (RA) as a repressor of both *Fgf8* domains and as an inducer of neural posteriorization in the trunk to allow hindbrain and spinal cord differentiation.

trunk development in a permissive fashion through its ability to repress caudal *Fgf8* expression to ensure proper spinal cord neuronal differentiation (Del Corral et al., 2003) and somitogenesis (Vermot et al., 2005; Sirbu and Duester, 2006) during body axis extension. A major target of RA repression during body axis extension has been found at the junction of the epiblast and neuroectoderm where RA restricts *Fgf8* expression to epiblast (primitive ectoderm), thus preventing it from extending anteriorly into node ectoderm and neuroectoderm (Sirbu and Duester, 2006). RA also represses caudal *Wnt8* expression, which plays a role in regulating *Raldh2* expression in presomitic mesoderm, suggesting that Wnt signaling may control the timing of the caudal RA-FGF differentiation switch (Olivera-Martinez and Storey, 2007). Additionally, RA represses *Fgf8* in the posterior mesoderm of the cardiac field (Ryckebusch et al., 2008; Sirbu et al., 2008). Thus, in the absence of RA signaling the epiblast *Fgf8* domain moves further to the anterior and the cardiac *Fgf8* domain moves further to the posterior. As RA signaling restricts the anteroposterior limits of each of the two early *Fgf8* expression domains (cardiac and epiblast), it creates an FGF8-free zone in between where the trunk develops (Figure 2). Overall, instructive and permissive actions of RA signaling during late gastrulation function to organize the embryonic trunk by allowing trunk progenitor cells to properly differentiate. Further evidence that RA functions as a trunk organizer has come from studies on embryos treated with excess RA, which transforms head structures such as forebrain/midbrain and anterior foregut endoderm to trunk neuroectoderm (hindbrain/spinal cord) and trunk foregut endoderm fates, respectively (Conlon and Rossant, 1992; Stafford and Prince, 2002).

As *Fgf8* is expressed prior to *Raldh2*, FGF8 signaling is already underway prior to the advent of RA signaling at E7.5 in mouse embryos (Sirbu et al., 2005). RA antagonism of FGF8 signaling along the anteroposterior axis then allows cells at the boundaries of these two signaling pathways to quickly transition away from FGF8 signaling and begin responding to RA signaling. *Fgf8* repression in the developing trunk during late gastrulation can thus be considered a fundamental permissive function of RA needed for trunk development, which is simultaneous to the well-established instructive role of RA as an inducer of homeobox genes in the posterior neuroectoderm and posterior foregut endoderm.

Tissues Derived from Neuroectoderm

The endogenous concentration of RA in mouse embryos is approximately 25 nM (Mic et al., 2003). Treatment of embryonic stem cells or embryonal carcinoma cells with levels of RA above 100 nM can induce a neural fate (Cai and Grabel, 2007), although studies on the normal function of RA in mouse embryos have demonstrated that neural induction does not require RA (Molotkova et al., 2005). The fact that RA is not synthesized in mouse embryos until E7.5 (well after induction of forebrain and midbrain neuroectoderm) is one indication that RA is not required for neural induction (Sirbu et al., 2005). Also, RA generated in the somitic mesoderm (E7.5–E8.5) stimulates RA signaling only in posterior neural tissues, and a loss of RA signaling does not affect expression of the neural induction markers *Sox1* and *Sox2* (Molotkova et al., 2005).

Instead, RA signaling acts upon neuroectoderm to influence its further differentiation. Investigation of embryos from single, double, and triple knockout mice lacking expression of *Raldh1*, *Raldh2*, and/or *Raldh3* indicates that the only action of RA required during early stages of neural development (up to E10.5) is in the posterior neuroectoderm (hindbrain and spinal cord) and eye (an outpocketing of the forebrain), but not in the forebrain itself (Molotkov et al., 2006; Molotkova et al., 2007). However, at later stages (after E12.5) *Raldh3* expression in the ventral forebrain is required for differentiation within the striatum, particularly for induction of dopamine receptor D2 in the nucleus accumbens (Molotkova et al., 2007).

RA treatment of amphibian embryos results in a loss of forebrain/midbrain accompanied by an anterior advance of hindbrain, suggesting that RA may be one of the factors regulating posterior transformation of the nervous system (Durstun et al., 1989; Sive et al., 1990). Indeed, vitamin A-deficient quail embryos and mouse embryos completely lacking RA signaling exhibit hindbrain defects, indicating that endogenous RA transforms the posterior neuroectoderm after neural induction occurs (Maden et al., 1996; Niederreither et al., 2000; Molotkova et al., 2005; Sirbu et al., 2005).

Hindbrain Patterning

Hox genes exhibit differential expression along the anteroposterior axis of the developing hindbrain and are intimately involved in rhombomere formation and identity (Krumlauf, 1993). A major function of RA signaling in the hindbrain involves control of *Hox* gene expression (Maden et al., 1996; Dupé and Lumsden, 2001). Several members of the *Hox* gene family are

direct targets of RA signaling including *Hoxb1* (Simeone et al., 1990), which is required for facial motor neuron differentiation in rhombomere 4 (r4). Prior to rhombomere formation, *Hoxb1* is expressed throughout the posterior hindbrain up to the presumptive r3/r4 border but soon becomes restricted to r4 (Wilkinson et al., 1989). *Hoxb1* is regulated by a retinoic acid response element (RARE) located 3' to the promoter. This element is required for early widespread induction in the posterior hindbrain up to the presumptive r3/4 boundary (Marshall et al., 1994). Interestingly, another RARE located 5' to the promoter is required for repression of *Hoxb1* in r3 and r5 to limit its expression to r4 (Studer et al., 1994). Additionally, repression of *Hoxb1* in r5 also depends upon the homeodomain protein encoded by *vHnf1* (*Hnf1b*), which is expressed in response to RA in the posterior hindbrain up to the r4/r5 boundary (Wiellette and Sive, 2003; Sirbu et al., 2005; Hernandez et al., 2004).

During establishment of *Hoxb1* expression, RA generated by RALDH2 in paraxial mesoderm initially travels as far anteriorly as presumptive r3 forming an early RA signaling boundary at r2/r3 just posterior to the forebrain/midbrain, which expresses the RA-degrading enzyme *Cyp26a1*. However, this boundary soon shifts posteriorly to the r4/r5 border due to expression of *Cyp26c1* in r4. Hence the hindbrain utilizes the RA-degrading functions of *Cyp26a1* and *Cyp26c1* to establish shifting boundaries of RA activity that induce both *Hoxb1* and *vHnf1*, a repressor of *Hoxb1* (Sirbu et al., 2005). Supporting this model, knockout and knockdown of *Cyp26a1* and *Cyp26c1* in mouse and zebrafish result in an anterior extension of *Hoxb1* expression into territory that normally develops into midbrain or forebrain (Hernandez et al., 2007; Uehara et al., 2007).

Motor Neuron Lineage Specification

In the spinal cord, sonic hedgehog (SHH) and RA are needed to establish a ventral fate that generates motor neurons (Sockanathan and Jessell, 1998; Del Corral et al., 2003; Novitch et al., 2003; Molotkova et al., 2005). Transcription factors required for motor neuron differentiation include Pax6 (expressed dorsally), Nkx6.1 (expressed ventrally), and Olig2 expressed in the region where Pax6 and Nkx6.1 overlap and where motor neurons develop (Marquardt and Pfaff, 2001). RA synthesized by RALDH2 in the adjacent somitic mesoderm travels to the spinal cord neuroectoderm. Mouse embryos lacking this source of RA fail to express Pax6 and Olig2 in the spinal cord, whereas Nkx6.1 expression is unaffected (Molotkova et al., 2005). In the absence of RA signaling, undifferentiated spinal cord neuroectoderm does not acquire a ventral motor neuron cell fate (Novitch et al., 2003; Molotkova et al., 2005). RA has been used as a differentiation agent along with SHH to generate motor neurons from both mouse (Wichterle et al., 2002) and human (Li et al., 2005) embryonic stem cells, providing a potential source of replacement cells for motor neuron diseases or spinal cord injuries.

Tissues Derived from Mesoderm

Bilateral Symmetry of Somites

Somitogenesis is the process whereby trunk paraxial mesoderm is sequentially segmented along the anteroposterior axis into bilaterally paired epithelial structures known as somites on the left and right sides of the embryonic axis. Rhythmic somite formation relies on a "clock and wavefront" mechanism in which

a molecular oscillator dependent upon Notch and Wnt signaling controls rhythmic expression of genes along the presomitic mesoderm (Pourquié, 2003). A moving wavefront of *Fgf8* gene expression in the primitive streak regresses posteriorly as the body axis extends, thus generating a somite determination front just anterior to the *Fgf8* expression domain (Dubrulle et al., 2001). A role for RA in somite development was suggested by studies showing that RA is required for the caudal expression of *Cdx1*, which is needed for proper development of the axial skeleton (Houle et al., 2003). The anteroposterior position of the determination front is also dependent upon RA generated in presomitic mesoderm, which represses caudal *Fgf8* expression (Del Corral et al., 2003).

Experiments in mouse, chick, and zebrafish embryos demonstrated that RALDH2 generates the RA required to maintain bilateral symmetry of the left and right columns of somites (Kawakami et al., 2005; Vermot et al., 2005; Vermot and Pourquié, 2005; Sirbu and Duester, 2006). Loss of RA signaling leads to a loss of left-right bilateral symmetry such that one side has fewer somites than the other. Presomitic mesoderm in RA-deficient embryos displays abnormal left-right asymmetric expression of *Hes7* and *Lfng*, which are required for Notch-dependent oscillator function during somitogenesis (Kawakami et al., 2005; Vermot et al., 2005; Vermot and Pourquié, 2005). Thus, a loss of RA allows left-right asymmetry to occur in presomitic mesoderm where it normally does not occur but does not alter left-right asymmetry normally observed in lateral plate mesoderm.

Expression of *Fgf8* mRNA in epiblast (primitive ectoderm) is shifted anteriorly in RALDH2-deficient embryos so that it enters the node ectoderm and neuroectoderm; thus it has been proposed that excessive anterior FGF signaling from ectoderm to mesoderm may be responsible for the somite defect (Sirbu and Duester, 2006). This hypothesis is supported by two observations. First, the somite determination front normally occurs in presomitic mesoderm just anterior to the *Fgf8* expression domain in the primitive streak, and an anterior shift in FGF signaling results in an anterior shift in somite position along the anteroposterior axis (Dubrulle et al., 2001). Second, as the FGF8 signal to the node is required for left-right asymmetry in lateral plate mesoderm (for instance, heart tube looping) (Meyers and Martin, 1999), it is possible that excessive FGF8 signaling to the node during RA deficiency may result in left-right asymmetry occurring also in the presomitic mesoderm where it should normally not occur.

Regulation of Heart Patterning

RA is required for anteroposterior patterning of the heart tube. Loss of RA synthesis leads to a severe reduction in the atria/inflow tract domain and the outflow tract/ventricular domain forms an abnormal cavity that is distended medially rather than undergoing rightward looping and septation into right and left ventricles (Niederreither et al., 2001). In both mouse and chick embryos, *Raldh2* is first expressed in the presomitic mesoderm, and then during the early stages of somitogenesis a caudorostral wave of *Raldh2* expression occurs in the lateral plate mesoderm up to a location just posterior to the cardiac crescent (Hochgreb et al., 2003). RA generated by RALDH2 travels into posterior heart mesoderm and is able to induce expression of an RA-reporter in the mouse heart from E7.5–E8.5.




Figure 3. Retinoic Acid Signaling during Early Heart Organogenesis

Retinoic acid (RA) generated by *retinaldehyde dehydrogenase 2* (*Raldh2*) in the somites and lateral plate mesoderm (lpm) travels anteriorly where it provides a signal that helps establish the posterior border of the heart field. The lateral domain of the heart field expresses *Tbx5* whereas the medial domain expresses *Isl1* as well as *Fgf8*, which is required for cardiac *Isl1* expression. Embryos from *Raldh2* knockout mice exhibit posterior expansion of *Fgf8* and *Isl1* into lateral plate mesoderm that normally is not part of the heart field. As the *Fgf8* promoter has been reported to contain an RA response element, RA may function as a repressor of cardiac *Fgf8*.

Studies in zebrafish have shown that RA restricts the size of the cardiac progenitor pool (Keegan et al., 2005), and studies in chick demonstrate that RA limits the location of ventricular progenitors along the anteroposterior axis (Hochgref et al., 2003). Analysis of cardiac genes in mouse embryos lacking RALDH2 suggests that the effect of RA on early heart development is not mediated through induction of target genes (Niederreither et al., 2001) but rather through repression of *Fgf8* expression in the posterior region of the heart (Ryckebusch et al., 2008; Sirbu et al., 2008).

During early heart tube organogenesis, studies in chick and mouse embryos have identified distinct progenitor cell populations in the splanchnic lateral plate mesoderm of the cardiac crescent that express unique combinations of transcription factors essential for heart development (Cai et al., 2003; Buckingham et al., 2005). For instance, *Tbx5* is expressed more laterally (sometimes referred to as first heart field), whereas the genes encoding transcription factor *Isl1* and the growth factor *Fgf8* are expressed more medially (sometimes referred to as second heart field). In mice, loss of RA signaling results in a loss of *Tbx5* expression in the posterior-lateral region of the cardiac crescent, but not the anterior region (Niederreither et al., 2001; Sirbu et al., 2008). This effect on *Tbx5* is indirect as RA signaling is required only in the posterior-medial domain where *Fgf8* and *Isl1* (but not *Tbx5*) are expressed (Sirbu et al., 2008). Mouse embryos lacking RALDH2 exhibit an increase in *Fgf8* and *Isl1* cardiac expression posteriorly (Ryckebusch et al., 2008; Sirbu et al., 2008) as well as an increase in FGF8 signaling marked by *Sprouty2* expression (Sirbu et al., 2008). Studies on embryos with cardiac-specific loss of *Fgf8* indicate that *Fgf8* is required for expression of *Isl1* in splanchnic mesodermal progenitors (Ilagan et al., 2006; Park et al., 2006). Due to the requirement of FGF8 signaling for *Isl1* induction, RA downregulation of *Fgf8* may thus normally limit the posterior extent of *Isl1* expression in cardiac mesoderm (Figure 3). Alternative promoters for *Fgf8* are controlled by a unique retinoic acid response element that allows expression of the major isoform (*Fgf8b*) when RAR is unliganded but represses *Fgf8b* when RA is present (Brondani et al., 2002). Thus, RA appears to function in a repressive rather than an inductive fashion during early cardiac organogenesis.

Forelimb Induction

Introduction of exogenous RA alters anteroposterior patterning of chick limb buds (Tickle et al., 1982) or proximodistal patterning of regenerating axolotl limbs (Maden, 1982). Later, it was demonstrated that *Shh* controls limb anteroposterior patterning and that RA-bead implants ectopically induce *Shh* (Riddle et al., 1993). Genetic studies in mice and zebrafish demonstrated that RA synthesis in the vicinity of limb buds is controlled by RALDH2 expressed in flank mesoderm lying next to the limb bud (Niederreither et al., 1999; Mic et al., 2002; Gibert et al., 2006). Further studies in mice demonstrated that RA activity in early limb buds is uniform across the anteroposterior axis but decreases from proximal to distal (Mic et al., 2004). It was proposed that opposing signals of RA, which is generated proximally in the flank, and *Fgf8*, which is expressed distally in the apical ectodermal ridge, may be important to control establishment of the limb proximodistal axis (Mercader et al., 2000). Whereas genetic support for *Fgf8* as a distal signal has been obtained, there is no genetic evidence that a proximal RA signaling center is required to establish the proximodistal axis of the limb (Tabin and Wolpert, 2007). In addition, studies on mouse embryos lacking the RA-degrading enzyme CYP26B1, which in wild-type embryos is expressed in the distal region of both forelimbs and hindlimbs, have shown that RA degradation is necessary for proper outgrowth of limb buds (Yashiro et al., 2004). Although loss of CYP26B1 results in an abnormal distal expansion of RA signaling in the forelimbs and hindlimbs that correlates with RA-induced limb teratogenesis, these findings do not provide evidence that a normal level of endogenous RA signaling is necessary for proximodistal patterning of the limbs.

In the absence of RA synthesis by RALDH2, forelimb buds do not develop and embryonic growth ceases prior to the stage when hindlimb buds are initiated (Niederreither et al., 2002; Mic et al., 2004); in zebrafish the absence of RA synthesis blocks induction of pectoral fin buds (Gibert et al., 2006). Thus, RA is required for induction of forelimb development, but whether RA also plays a later role in patterning of the limbs remains controversial. Rescue of *Raldh2* mutant embryos by low-dose maternal dietary RA supplementation rescues lethality and results in limb bud induction despite no detectable RA activity in forelimbs or hindlimbs; in this model, forelimb buds

are undersized whereas hindlimb buds appear normal (Niederreither et al., 2002; Mic et al., 2004), suggesting that RA may not have a role in hindlimb induction and patterning. Further support for this conclusion has come from *Rdh10* mutant embryos (lacking an enzyme upstream of RALDH2 for RA synthesis), which survive long enough to exhibit small forelimbs and normal hindlimbs similar to RA-rescued *Raldh2* mutants (Sandell et al., 2007). However, hindlimb buds may be affected by RA signaling that occurs in the mesonephros that expresses *Raldh3* (Mic et al., 2002). Thus, studies examining whether low-dose dietary RA can rescue limb development in embryos lacking both RALDH2 and RALDH3 will be needed to assess whether hindlimbs require RA for induction or for patterning along either the proximodistal or anteroposterior axes.

The rescue of *Raldh2* null mutant embryos with low-dose maternal dietary supplementation of RA provides less RA than normally synthesized by RALDH2 (Mic et al., 2003), resulting in conditions where RA activity is absent in forelimb mesoderm but present nearby in the body axis, particularly the neuroectoderm (Niederreither et al., 2002; Mic et al., 2004); neuroectoderm expresses very high levels of CRABP2 and RAR β and can presumably respond to lower RA levels than somitic and limb mesoderm. This observation suggests that although RA signaling is unnecessary in forelimb mesoderm for forelimb budding, RA may act in the neuroectoderm to permit limb induction. Given that neuroectoderm is unnecessary for induction of forelimbs (Rong et al., 1992), RA activity in neuroectoderm may instead downregulate expression of a diffusible factor in the body axis that inhibits forelimb induction. Three observations suggest that this factor, if it exists, could be FGF8. (1) RA represses caudal *Fgf8* expression during body axis extension (Del Corral et al., 2003; Vermot et al., 2005); (2) ectopic FGF signaling from beads implanted in the chick wing field impairs wing development (Cohn et al., 1995; Mercader et al., 2000); (3) RA repression of *Fgf8* expression during mouse body axis extension occurs at the junction of the neuroectoderm/epiblast during the somite stages when forelimb induction occurs (Sirbu and Duester, 2006). Thus, rather than playing an instructive role in forelimb budding, recent findings suggest that RA generated in somites may play a permissive role in forelimb induction through its action in the body axis near the forelimb field. At later steps when limb patterning occurs, it is doubtful that RA has a role.

Organs Derived from Endoderm

Pancreas Induction

Many studies of pancreas development have focused on the homeobox transcription factor *Pdx1*, which is required for pancreatic specification in the posterior foregut. *Pdx1* is highly expressed in the dorsal and ventral endodermal buds that give rise to the mature pancreas. Evidence suggests that RA may be the mesodermal signal required for initiation of *Pdx1* expression (Stafford and Prince, 2002). Zebrafish embryos deficient in RA lack expression of *Pdx1* and consequently fail to induce pancreas development. Similar results have also been obtained in mouse (Martín et al., 2005; Molotkov et al., 2005) and frog (Chen et al., 2004) embryos. Studies on avian embryos revealed that signals from the lateral plate mesoderm can drive endodermal cells to express *Pdx1* and thus generate

a pancreatic fate (Kumar et al., 2003). Examination of mouse embryos carrying the RA-reporter transgene demonstrated that the pancreas is normally exposed to RA generated in the surrounding splanchnic lateral plate mesoderm (Molotkov et al., 2005).

In RALDH2-deficient mouse embryos (Molotkov et al., 2005) and RAR antagonist-treated *Xenopus* embryos (Chen et al., 2004), dorsal endodermal pancreatic tissue is not correctly specified. In contrast, ventral endodermal pancreatic tissue and liver are still specified, demonstrating that only dorsal endoderm requires RA activity for pancreatic development. Specification of dorsal pancreatic tissue can be rescued in RALDH2-deficient embryos by low-dose maternal administration of RA, a treatment that preferentially restores RA activity to the dorsal endoderm but not the surrounding dorsal mesoderm (Molotkov et al., 2005). This suggests that RA acts directly in the dorsal endoderm for pancreas specification rather than in the lateral plate mesoderm where RA is synthesized. In zebrafish, transplantation studies have shown that RAR needs to act only in the endoderm for pancreas induction and that the RA is derived from the surrounding mesoderm (Stafford et al., 2006).

Lung Induction

The lung is another organ derived from the endoderm of the posterior foregut that requires RA signaling for organogenesis. RA from the splanchnic mesoderm surrounding the endoderm has been found to be important for stimulation of posterior foregut endoderm to a lung fate at E9.5 in the mouse (Malpel et al., 2000). RALDH2-deficient embryos rescued from early lethality by maternal dietary RA between E7.5 and E8.5 fail to develop lungs and lack RA signaling in the foregut (Wang et al., 2006). In RA-deficient embryos the primary lung bud is specified, but it does not express the RA-inducible *Hoxa5* gene and it is unable to achieve outgrowth or branching due to a loss of both *Fgf10* expression and FGF10 signaling in the lung epithelium; treatment of RA-deficient embryos in vitro with FGF10 can restore lung budding and branching (Wang et al., 2006). Further studies demonstrated that a loss of RA signaling in the foregut results in upregulation of TGF- β 1 and TGF- β target genes, and treatment of wild-type embryos with exogenous TGF- β 1 can reproduce the lung bud defect seen in RA-deficient embryos (Chen et al., 2007). Therefore, it appears that RA functions during lung budding both as an inducer of *Hoxa5* and an inhibitor of TGF- β 1 signaling, which then permits local expression of *Fgf10* needed for expansion of the lung bud and branching.

Morphogenetic Movements during Eye Formation

As all three mouse *Raldh* genes are expressed in the developing eye, it is of interest to discuss in more detail their expression patterns and what their gene knockouts have revealed so far about the mechanism of RA signaling during eye development. Vitamin A deficiency and reduced RA receptor function in RAR null mice (Lohnes et al., 1994) both result in incomplete closure of the choroid fissure (ocular coloboma) as well as microphthalmia (small eyes) and abnormalities of the cornea, eyelids, and conjunctiva observed during late fetal stages. However, due to remaining RA activity these studies could not determine at what developmental stage(s) RA is required for eye development.




Figure 4. Retinoic Acid Regulates Eye Development

Paracrine retinoic acid (RA) signaling controls two distinct phases of eye morphogenetic movements and involves all three *retinaldehyde dehydrogenase* (*Raldh*) genes. During invagination of the optic vesicle to form an optic cup (E9.5–E10.5 in mouse), RA generated by *Raldh2* in the perioptic mesenchyme as well as *Raldh3* in tissue fated to become the dorsal retinal pigment epithelium (RPE) travels to the neural retina where it is required for ventral invagination during optic cup formation (E10.5–birth). After optic cup formation (E10.5–birth), *Raldh2* is no longer expressed in the perioptic mesenchyme and *Raldh3* expression ends in the RPE, but *Raldh1* and *Raldh3* are now expressed in the dorsal and ventral neural retina, respectively. RA generated by *Raldh1* and *Raldh3* is not required for patterning the neural retina, but this RA travels outside the retina where it limits invasion of perioptic mesenchyme during anterior eye development (cornea and eyelid formation).

The three RALDHs are expressed in distinct tissues during development of the mouse eye, and in each case RA signaling activity has been detected in nearby tissues and found to be required for eye development (Molotkov et al., 2006). Invagination of the optic vesicle epithelium from E9.5–E10.5 results in formation of an optic cup in which the epithelium has folded to form separate layers for neural retina and retinal pigment epithelium, with both being folded around the lens vesicle that has developed by invagination of the surface ectoderm (Figure 4). RALDH2 generates RA at E9.0 in the perioptic mesenchyme next to the temporal side of the optic vesicle whereas RALDH3 generates RA in the dorsal retinal pigment epithelium starting at E9.5 just prior to invagination; a lack of both RALDH2 and RALDH3 results in a failure of ventral optic cup invagination (Molotkov et al., 2006). During optic cup formation the following changes in *Raldh* function occur: (1) *Raldh2* expression near the optic cup terminates at E10.0; (2) *Raldh3* expression terminates in the dorsal retinal pigment epithelium and initiates in the ventral neural retina at E10.5; (3) RALDH1 begins to generate RA in the dorsal neural retina at E10.5. At E11.5 when the ventral portion of the optic cup forms the choroid fissure, expression of *Raldh1* and *Raldh3* continues in the dorsal and ventral neural retina, respectively, and endures even after closure of the optic fissure at E13.5. However, RALDH1 and RALDH3 do not generate RA for retinal dorsoventral patterning, but instead they have completely redundant functions in generating RA that travels from the retina to the perioptic mesenchyme where it stimulates apoptosis; a lack of both RALDH1 and RALDH3 results in mesenchymal overgrowth in the cornea and eyelids (Matt et al., 2005; Molotkov et al., 2006). *Pitx2* is a potential target of ocular RA signaling as its expression is severely downregulated in perioptic mesenchyme of mutant embryos lacking both RALDH1 and RALDH3 (Matt et al., 2005).

These findings suggest the existence of two distinct phases of RA signaling required for eye development: an early phase for optic cup formation and a late phase for anterior eye formation (Figure 4). In both cases, cells expressing *Raldh* genes provide an RA signal that functions to control morphogenetic movements in neighboring cells. Also, in both cases two *Raldh* genes function as RA sources, providing functional redundancy. The target of RA action changes dur-

ing eye morphogenesis. The initial target is the invaginating neural retina at the optic vesicle stage with the sources of RA being *Raldh2* expressed in mesenchyme located temporally to the optic vesicle and *Raldh3* expressed in the retinal pigment epithelium. After optic cup formation the target switches to the perioptic mesenchyme with the sources being *Raldh1* expressed in the dorsal retina and *Raldh3* expressed in the ventral retina. These targets are distinct from but adjacent to locations of RA synthesis, thus demonstrating that RA functions in a paracrine fashion to guide morphogenetic movements of neighboring cells.

Development of the Genitourinary Tract

RA controls some aspects of genitourinary tract development as *Raldh2* is expressed in mesenchymal cells of the mesonephros and stromal cells of the developing kidney, and *Raldh3* is expressed in the ureteric bud (Batourina et al., 2001; Mic et al., 2002). Through analysis of RA receptor knockout mice, RA signaling has been found to play a key role in controlling epithelial/mesenchymal interactions during kidney development through induction of *Ret* expression (Batourina et al., 2001). Also, RA generated in the urogenital sinus stimulates apoptosis in the common nephric duct needed for establishing the connections between the ureters and bladder (Batourina et al., 2005).

RA has recently been found to stimulate sex-specific onset of meiosis in germ cells of mice. The onset of meiosis occurs earlier in the ovary (E13.5) than in the testis (after birth) (Bowles et al., 2006; Koubova et al., 2006). RA (likely from the mesonephros) induces expression of *Stra8* in germ cells, which is needed for the transition into meiosis in both the ovary and testis. In the testis the RA-degrading enzyme CYP26B1 is expressed and prevents induction of *Stra8* expression. Premature onset of meiosis in the testis occurs in mice lacking CYP26B1 (Bowles et al., 2006) and in organ cultures treated with a CYP26 inhibitor (Koubova et al., 2006).

Paracrine Signaling by Retinoic Acid

In summary, there are many examples of paracrine RA signaling (Table 3). The existence of autocrine RA signaling is not yet ruled out, but further studies will be needed to determine if this actually occurs. Future investigations of this type should pro-

Table 3. Examples of the Paracrine Function of Retinoic Acid

Retinoic Acid Source	Retinoic Acid Target
Somitic mesoderm (<i>Raldh2</i>)	Hindbrain neuroectoderm (anteroposterior patterning)
Somitic mesoderm (<i>Raldh2</i>)	Spinal cord neuroectoderm (motor neuron formation)
Somitic mesoderm (<i>Raldh2</i>)	Posterior neuroectoderm & node ectoderm (somitogenesis)
Somitic mesoderm (<i>Raldh2</i>)	Posterior body axis (forelimb induction)
Somitic/lateral plate mesoderm (<i>Raldh2</i>)	Cardiac mesoderm (anteroposterior patterning)
Lateral plate mesoderm (<i>Raldh2</i>)	Foregut endoderm (pancreas & lung induction)
Optic vesicle mesenchyme (<i>Raldh2</i>)	Neural retina (ventral optic cup formation)
Retinal pigment epithelium (<i>Raldh3</i>)	Neural retina (ventral optic cup formation)
Neural retina (<i>Raldh1</i> and <i>Raldh3</i>)	Periopic mesenchyme (eyelid and cornea development)
Mesonephros (<i>Raldh2</i>)	Ureteric bud (kidney induction)
Mesonephros (<i>Raldh2</i>)	Ovary (meiotic induction)

vide a more complete understanding of when and where RA is generated in embryos, how far it can travel from its site of synthesis, what the target tissues are, what developmental processes it affects, and what genes it regulates. Such knowledge will be essential for understanding mammalian organogenesis and will facilitate the development of rational strategies for optimal use of RA with other reagents to reliably differentiate stem cells into specific cell types that can be used to develop stem cell-based treatments for disease.

ACKNOWLEDGMENTS

I am grateful to past and present members of my laboratory for provocative discussions. Funding was provided by NIH grants GM062848 and EY013969 and California Institute of Regenerative Medicine grant RS1-00193.

REFERENCES

- Abu-Abed, S., Dollé, P., Metzger, D., Beckett, B., Chambon, P., and Petkovich, M. (2001). The retinoic acid-metabolizing enzyme, CYP26A1, is essential for normal hindbrain patterning, vertebral identity, and development of posterior structures. *Genes Dev.* 15, 226–240.
- Ang, H.L., Deltour, L., Hayamizu, T.F., Zgombic-Knight, M., and Duester, G. (1996). Retinoic acid synthesis in mouse embryos during gastrulation and craniofacial development linked to class IV alcohol dehydrogenase gene expression. *J. Biol. Chem.* 271, 9526–9534.
- Arnhold, T., Tzimas, G., Wittfoht, W., Plonait, S., and Nau, H. (1996). Identification of 9-cis-retinoic acid, 9,13-di-cis-retinoic acid, and 14-hydroxy-4,14-retro-retinol in human plasma after liver consumption. *Life Sci.* 59, PL 169–PL 177.
- Batourina, E., Gim, S., Bello, N., Shy, M., Clagett-Dame, M., Srinivas, S., Costantini, F., and Mendelsohn, C. (2001). Vitamin A controls epithelial/mesenchymal interactions through *Ref* expression. *Nat. Genet.* 27, 74–78.
- Batourina, E., Tsai, S., Lambert, S., Sprengle, P., Viana, R., Dutta, S., Hensle, T., Wang, F.W., Niederreither, K., McMahon, A.P., et al. (2005). Apoptosis induced by vitamin A signaling is crucial for connecting the ureters to the bladder. *Nat. Genet.* 37, 1082–1089.
- Bowles, J., Knight, D., Smith, C., Wilhelm, D., Richman, J., Mamiya, S., Yashiro, K., Chawengsaksophak, K., Wilson, M.J., Rossant, J., et al. (2006). Retinoid signaling determines germ cell fate in mice. *Science* 312, 596–600.
- Brondani, V., Klimkait, T., Egly, J.M., and Hamy, F. (2002). Promoter of FGF8 reveals a unique regulation by unliganded RARα. *J. Mol. Biol.* 319, 715–728.
- Buckingham, M., Meilhac, S., and Zaffran, S. (2005). Building the mammalian heart from two sources of myocardial cells. *Nat. Rev. Genet.* 6, 826–835.
- Cai, C., and Grabel, L. (2007). Directing the differentiation of embryonic stem cells to neural stem cells. *Dev. Dyn.* 236, 3255–3266.
- Cai, C.L., Liang, X., Shi, Y., Chu, P.H., Pfaff, S.L., Chen, J., and Evans, S. (2003). Isl1 identifies a cardiac progenitor population that proliferates prior to differentiation and contributes a majority of cells to the heart. *Dev. Cell* 5, 877–889.
- Chawla, A., Repa, J.J., Evans, R.M., and Mangelsdorf, D.J. (2001). Nuclear receptors and lipid physiology: Opening the X-files. *Science* 294, 1866–1870.
- Chen, F., Desai, T.J., Qian, J., Niederreither, K., Lu, J., and Cardoso, W.V. (2007). Inhibition of Tgfb signaling by endogenous retinoic acid is essential for primary lung bud induction. *Development* 134, 2969–2979.
- Chen, Y., Pan, F.C., Brandes, N., Afelik, S., Solter, M., and Pieler, T. (2004). Retinoic acid signaling is essential for pancreas development and promotes endocrine at the expense of exocrine cell differentiation in *Xenopus*. *Dev. Biol.* 271, 144–160.
- Clagett-Dame, M., and DeLuca, H.F. (2002). The role of vitamin A in mammalian reproduction and embryonic development. *Annu. Rev. Nutr.* 22, 347–381.
- Cohn, M.J., Izpisua-Belmonte, J.C., Abud, H., Heath, J.K., and Tickle, C. (1995). Fibroblast growth factors induce additional limb development from the flank of chick embryos. *Cell* 80, 739–746.
- Conlon, R.A., and Rossant, J. (1992). Exogenous retinoic acid rapidly induces anterior ectopic expression of murine *Hox-2* genes in vivo. *Development* 116, 357–368.
- Del Corral, R.D., Olivera-Martinez, I., Goriely, A., Gale, E., Maden, M., and Storey, K. (2003). Opposing FGF and retinoid pathways control ventral neural pattern, neuronal differentiation, and segmentation during body axis extension. *Neuron* 40, 65–79.
- Dersch, H., and Zile, M.H. (1993). Induction of normal cardiovascular development in the vitamin A-deprived quail embryo by natural retinoids. *Dev. Biol.* 160, 424–433.
- Dickman, E.D., Thaller, C., and Smith, S.M. (1997). Temporally-regulated retinoic acid depletion produces specific neural crest, ocular and nervous system defects. *Development* 124, 3111–3121.
- Dubrulle, J., McGrew, M.J., and Pourquie, O. (2001). FGF signaling controls somite boundary position and regulates segmentation clock control of spatiotemporal *Hox* gene activation. *Cell* 106, 219–232.
- Dupé, V., and Lumsden, A. (2001). Hindbrain patterning involves graded responses to retinoic acid signalling. *Development* 128, 2199–2208.
- Durston, A.J., Timmermans, J.P.M., Hage, W.J., Hendriks, H.F.J., De Vries, N.J., Heideveld, M., and Nieuwkoop, P.D. (1989). Retinoic acid causes an anteroposterior transformation in the developing central nervous system. *Nature* 340, 140–144.
- Gallego, O., Belyaeva, O.V., Porte, S., Ruiz, F.X., Stetsenko, A.V., Shabrova, E.V.,

- Kostereva, N.V., Farres, J., Pares, X., and Kedishvili, N.Y. (2006). Comparative functional analysis of human medium-chain dehydrogenases, short-chain dehydrogenases/reductases and aldo-keto reductases with retinoids. *Biochem. J.* 399, 101–109.
- Germain, P., Iyer, J., Zechel, C., and Gronemeyer, H. (2002). Co-regulator recruitment and the mechanism of retinoic acid receptor synergy. *Nature* 415, 187–192.
- Ghyselinck, N.B., Båvik, C., Sapin, V., Mark, M., Bonnier, D., Hindelang, C., Dierich, A., Nilsson, C.B., Håkansson, H., Sauvaut, P., et al. (1999). Cellular retinol-binding protein I is essential for vitamin A homeostasis. *EMBO J.* 18, 4903–4914.
- Gibert, Y., Gajewski, A., Meyer, A., and Begemann, G. (2006). Induction and prepatterning of the zebrafish pectoral fin bud requires axial retinoic acid signaling. *Development* 133, 2649–2659.
- Hernandez, R.E., Rikhof, H.A., Bachmann, R., and Moens, C.B. (2004). *vhnf1* integrates global RA patterning and local FGF signals to direct posterior hind-brain development in zebrafish. *Development* 131, 4511–4520.
- Hernandez, R.E., Putzke, A.P., Myers, J.P., Margaretha, L., and Moens, C.B. (2007). Cyp26 enzymes generate the retinoic acid response pattern necessary for hindbrain development. *Development* 134, 177–187.
- Hochgreb, T., Linhares, V.L., Menezes, D.C., Sampaio, A.C., Yan, C.Y.I., Cardoso, W.V., Rosenthal, N., and Xavier-Neto, J. (2003). A caudorostral wave of RALDH2 conveys anteroposterior information to the cardiac field. *Development* 130, 5363–5374.
- Houle, M., Sylvestre, J.R., and Lohnes, D. (2003). Retinoic acid regulates a subset of Cdx1 function in vivo. *Development* 130, 6555–6567.
- Ilagan, R., Abu-Issa, R., Brown, D., Yang, Y.P., Jiao, K., Schwartz, R.J., Klingensmith, J., and Meyers, E.N. (2006). Fgf8 is required for anterior heart field development. *Development* 133, 2435–2445.
- Kawaguchi, R., Yu, J., Honda, J., Hu, J., Whitelegge, J., Ping, P., Wiita, P., Bok, D., and Sun, H. (2007). A membrane receptor for retinol binding protein mediates cellular uptake of vitamin A. *Science* 315, 820–825.
- Kawakami, Y., Raya, A., Raya, R.M., Rodriguez-Esteban, C., and Izpisua-Belmonte, J.C. (2005). Retinoic acid signalling links left-right asymmetric patterning and bilaterally symmetric somitogenesis in the zebrafish embryo. *Nature* 435, 165–171.
- Keegan, B.R., Feldman, J.L., Begemann, G., Ingham, P.W., and Yelon, D. (2005). Retinoic acid signaling restricts the cardiac progenitor pool. *Science* 307, 247–249.
- Koubova, J., Menke, D.B., Zhou, Q., Capel, B., Griswold, M.D., and Page, D.C. (2006). Retinoic acid regulates sex-specific timing of meiotic initiation in mice. *Proc. Natl. Acad. Sci. USA* 103, 2474–2479.
- Krumlauf, R. (1993). *Hox* genes and pattern formation in the branchial region of the vertebrate head. *Trends Genet.* 9, 106–112.
- Kumar, M., Jordan, N., Melton, D., and Grapin-Botton, A. (2003). Signals from lateral plate mesoderm instruct endoderm toward a pancreatic fate. *Dev. Biol.* 259, 109–122.
- Li, X.J., Du, Z.W., Zarnowska, E.D., Pankratz, M., Hansen, L.O., Pearce, R.A., and Zhang, S.C. (2005). Specification of motoneurons from human embryonic stem cells. *Nat. Biotechnol.* 23, 215–221.
- Lohnes, D., Mark, M., Mendelsohn, C., Dollé, P., Dierich, A., Gorry, P., Gansmuller, A., and Chambon, P. (1994). Function of the retinoic acid receptors (RARs) during development. (I) Craniofacial and skeletal abnormalities in RAR double mutants. *Development* 120, 2723–2748.
- Maden, M. (1982). Vitamin A and pattern formation in the regenerating limb. *Nature* 295, 672–675.
- Maden, M., Gale, E., Kostetskii, I., and Zile, M.H. (1996). Vitamin A-deficient quail embryos have half a hindbrain and other neural defects. *Curr. Biol.* 6, 417–426.
- Malpel, S., Mendelsohn, C., and Cardoso, W.V. (2000). Regulation of retinoic acid signaling during lung morphogenesis. *Development* 127, 3057–3067.
- Marlétaz, F., Holland, L.Z., Laudet, V., and Schubert, M. (2006). Retinoic acid signaling and the evolution of chordates. *Int. J. Biol. Sci.* 2, 38–47.
- Marquardt, T., and Pfaff, S.L. (2001). Cracking the transcriptional code for cell specification in the neural tube. *Cell* 106, 651–654.
- Marshall, H., Studer, M., Pöpperl, H., Aparicio, S., Kuroiwa, A., Brenner, S., and Krumlauf, R. (1994). A conserved retinoic acid response element required for early expression of the homeobox gene *Hoxb-1*. *Nature* 370, 567–571.
- Martín, M., Gallego-Llamas, J., Ribes, V., Keding, M., Niederreither, K., Chambon, P., Dollé, P., and Gradwohl, G. (2005). Dorsal pancreas agenesis in retinoic acid-deficient *Raldh2* mutant mice. *Dev. Biol.* 284, 399–411.
- Matt, N., Dupé, V., Garnier, J.-M., Dennefeld, C., Chambon, P., Mark, M., and Ghyselinck, N.B. (2005). Retinoic acid-dependent eye morphogenesis is orchestrated by neural crest cells. *Development* 132, 4789–4800.
- Mendelsohn, C., Lohnes, D., Decimo, D., Lufkin, T., Le Meur, M., Chambon, P., and Mark, M. (1994). Function of the retinoic acid receptors (RARs) during development. (II) Multiple abnormalities at various stages of organogenesis in RAR double mutants. *Development* 120, 2749–2771.
- Mercader, N., Leonardo, E., Piedra, M.E., Martínez-A, C., Ros, M.A., and Torres, M. (2000). Opposing RA and FGF signals control proximodistal vertebrate limb development through regulation of Meis genes. *Development* 127, 3961–3970.
- Meyers, E.N., and Martin, G.R. (1999). Differences in left-right axis pathways in mouse and chick: functions of FGF8 and SHH. *Science* 285, 403–406.
- Mic, F.A., Haselbeck, R.J., Cuenca, A.E., and Duester, G. (2002). Novel retinoic acid generating activities in the neural tube and heart identified by conditional rescue of *Raldh2* null mutant mice. *Development* 129, 2271–2282.
- Mic, F.A., Molotkov, A., Benbrook, D.M., and Duester, G. (2003). Retinoid activation of retinoic acid receptor but not retinoid X receptor is sufficient to rescue lethal defect in retinoic acid synthesis. *Proc. Natl. Acad. Sci. USA* 100, 7135–7140.
- Mic, F.A., Sirbu, I.O., and Duester, G. (2004). Retinoic acid synthesis controlled by *Raldh2* is required early for limb bud initiation and then later as a proximodistal signal during apical ectodermal ridge formation. *J. Biol. Chem.* 279, 26698–26706.
- Molotkov, A., Deltour, L., Foglio, M.H., Cuenca, A.E., and Duester, G. (2002a). Distinct retinoid metabolic functions for alcohol dehydrogenase genes *Adh1* and *Adh4* in protection against vitamin A toxicity or deficiency revealed in double null mutant mice. *J. Biol. Chem.* 277, 13804–13811.
- Molotkov, A., Fan, X., Deltour, L., Foglio, M.H., Martras, S., Farrés, J., Parés, X., and Duester, G. (2002b). Stimulation of retinoic acid production and growth by ubiquitously-expressed alcohol dehydrogenase *Adh3*. *Proc. Natl. Acad. Sci. USA* 99, 5337–5342.
- Molotkov, A., Ghyselinck, N.B., Chambon, P., and Duester, G. (2004). Opposing actions of cellular retinol-binding protein and alcohol dehydrogenase control the balance between retinol storage and degradation. *Biochem. J.* 383, 295–302.
- Molotkov, A., Molotkova, N., and Duester, G. (2005). Retinoic acid generated by *Raldh2* in mesoderm is required for mouse dorsal endodermal pancreas development. *Dev. Dyn.* 232, 950–957.
- Molotkov, A., Molotkova, N., and Duester, G. (2006). Retinoic acid guides eye morphogenetic movements via paracrine signaling but is unnecessary for retinal dorsoventral patterning. *Development* 133, 1901–1910.
- Molotkova, N., Molotkov, A., Sirbu, I.O., and Duester, G. (2005). Requirement of mesodermal retinoic acid generated by *Raldh2* for posterior neural transformation. *Mech. Dev.* 122, 145–155.
- Molotkova, N., Molotkov, A., and Duester, G. (2007). Role of retinoic acid during forebrain development begins late when *Raldh3* generates retinoic acid in the ventral subventricular zone. *Dev. Biol.* 303, 601–610.
- Niederreither, K., Subbarayan, V., Dollé, P., and Chambon, P. (1999).

- Embryonic retinoic acid synthesis is essential for early mouse post-implantation development. *Nat. Genet.* 21, 444–448.
- Niederreither, K., Vermot, J., Schuhbauer, B., Chambon, P., and Dollé, P. (2000). Retinoic acid synthesis and hindbrain patterning in the mouse embryo. *Development* 127, 75–85.
- Niederreither, K., Vermot, J., Messaddeq, N., Schuhbauer, B., Chambon, P., and Dollé, P. (2001). Embryonic retinoic acid synthesis is essential for heart morphogenesis in the mouse. *Development* 128, 1019–1031.
- Niederreither, K., Vermot, J., Schuhbauer, B., Chambon, P., and Dollé, P. (2002). Embryonic retinoic acid synthesis is required for forelimb growth and anteroposterior patterning in the mouse. *Development* 129, 3563–3574.
- Novitsch, B.G., Wichterle, H., Jessell, T.M., and Sockanathan, S. (2003). A requirement for retinoic acid-mediated transcriptional activation in ventral neural patterning and motor neuron specification. *Neuron* 40, 81–95.
- Noy, N. (2000). Retinoid-binding proteins: mediators of retinoid action. *Biochem. J.* 348, 481–495.
- Olivera-Martinez, I., and Storey, K.G. (2007). Wnt signals provide a timing mechanism for the FGF-retinoid differentiation switch during vertebrate body axis extension. *Development* 134, 2125–2135.
- Park, E.J., Ogden, L.A., Talbot, A., Evans, S., Cai, C.L., Black, B.L., Frank, D.U., and Moon, A.M. (2006). Required, tissue-specific roles for Fgf8 in outflow tract formation and remodeling. *Development* 133, 2419–2433.
- Pourquié, O. (2003). The segmentation clock: converting embryonic time into spatial pattern. *Science* 301, 328–330.
- Quadro, L., Blaner, W.S., Salchow, D.J., Vogel, S., Piantadosi, R., Gouras, P., Freeman, S., Cosma, M.P., Colantuoni, V., and Gottesman, M.E. (1999). Impaired retinal function and vitamin A availability in mice lacking retinol-binding protein. *EMBO J.* 18, 4633–4644.
- Riddle, R.D., Johnson, R.L., Laufer, E., and Tabin, C. (1993). *Sonic hedgehog* mediates the polarizing activity of the ZPA. *Cell* 75, 1401–1416.
- Rong, P.M., Teillet, M.-A., Ziller, C., and Le Douarin, N.M. (1992). The neural tube/notochord complex is necessary for vertebral but not limb and body wall striated muscle differentiation. *Development* 115, 657–672.
- Rossant, J., Zirngibl, R., Cado, D., Shago, M., and Giguère, V. (1991). Expression of a retinoic acid response element-*hsp*lacZ transgene defines specific domains of transcriptional activity during mouse embryogenesis. *Genes Dev.* 5, 1333–1344.
- Ryckebusch, L., Wang, Z., Bertrand, N., Lin, S.-C., Chi, X., Schwartz, R., Zafra, S., and Niederreither, K. (2008). Retinoic acid deficiency alters second heart field formation. *Proc. Natl. Acad. Sci. USA* 105, 2913–2918.
- Sandell, L.L., Sanderson, B.W., Moiseyev, G., Johnson, T., Mushegian, A., Young, K., Rey, J.P., Ma, J.X., Staehling-Hampton, K., and Trainor, P.A. (2007). RDH10 is essential for synthesis of embryonic retinoic acid and is required for limb, craniofacial, and organ development. *Genes Dev.* 21, 1113–1124.
- Schug, T.T., Berry, D.C., Shaw, N.S., Travis, S.N., and Noy, N. (2007). Opposing effects of retinoic acid on cell growth result from alternate activation of two different nuclear receptors. *Cell* 129, 723–733.
- Sessler, R.J., and Noy, N. (2005). A ligand-activated nuclear localization signal in cellular retinoic acid binding protein-II. *Mol. Cell* 18, 343–353.
- Simeone, A., Acampora, D., Arcioni, L., Andrews, P.W., Boncinelli, E., and Mavilio, F. (1990). Sequential activation of *HOX2* homeobox genes by retinoic acid in human embryonal carcinoma cells. *Nature* 346, 763–766.
- Sirbu, I.O., and Duester, G. (2006). Retinoic acid signaling in node ectoderm and posterior neural plate directs left-right patterning of somitic mesoderm. *Nat. Cell Biol.* 8, 271–277.
- Sirbu, I.O., Gresh, L., Barra, J., and Duester, G. (2005). Shifting boundaries of retinoic acid activity control hindbrain segmental gene expression. *Development* 132, 2611–2622.
- Sirbu, I.O., Zhao, X., and Duester, G. (2008). Retinoic acid controls heart anteroposterior patterning by down-regulating *Isl1* through the *Fgf8* pathway. *Dev. Dyn.* 237, 1627–1635.
- Sive, H.L., Draper, B.W., Harland, R.M., and Weintraub, H. (1990). Identification of a retinoic acid-sensitive period during primary axis formation in *Xenopus laevis*. *Genes Dev.* 4, 932–942.
- Sockanathan, S., and Jessell, T.M. (1998). Motor neuron-derived retinoid signaling specifies the subtype identity of spinal motor neurons. *Cell* 94, 503–514.
- Stafford, D., and Prince, V.E. (2002). Retinoic acid signaling is required for a critical early step in zebrafish pancreatic development. *Curr. Biol.* 12, 1215–1220.
- Stafford, D., White, R.J., Kinkel, M.D., Linville, A., Schilling, T.F., and Prince, V.E. (2006). Retinoids signal directly to zebrafish endoderm to specify insulin-expressing beta-cells. *Development* 133, 949–956.
- Studer, M., Pöpperl, H., Marshall, H., Kuroiwa, A., and Krumlauf, R. (1994). Role of a conserved retinoic acid response element in rhombomere restriction of *Hoxb-1*. *Science* 265, 1728–1732.
- Tabin, C., and Wolpert, L. (2007). Rethinking the proximodistal axis of the vertebrate limb in the molecular era. *Genes Dev.* 21, 1433–1442.
- Tickle, C., Alberts, B.M., Wolpert, L., and Lee, J. (1982). Local application of retinoic acid to the limb bud mimics the action of the polarizing region. *Nature* 296, 564–565.
- Uehara, M., Yashiro, K., Mamiya, S., Nishino, J., Chambon, P., Dolle, P., and Sakai, Y. (2007). CYP26A1 and CYP26C1 cooperatively regulate anterior-posterior patterning of the developing brain and the production of migratory cranial neural crest cells in the mouse. *Dev. Biol.* 302, 399–411.
- Vermot, J., and Pourquié, O. (2005). Retinoic acid coordinates somitogenesis and left-right patterning in vertebrate embryos. *Nature* 435, 215–220.
- Vermot, J., Llamas, J.G., Fraulob, V., Niederreither, K., Chambon, P., and Dollé, P. (2005). Retinoic acid controls the bilateral symmetry of somite formation in the mouse embryo. *Science* 308, 563–566.
- Wang, Z., Dolle, P., Cardoso, W.V., and Niederreither, K. (2006). Retinoic acid regulates morphogenesis and patterning of posterior foregut derivatives. *Dev. Biol.* 297, 433–445.
- Wichterle, H., Lieberam, I., Porter, J.A., and Jessell, T.M. (2002). Directed differentiation of embryonic stem cells into motor neurons. *Cell* 110, 385–397.
- Wiellette, E.L., and Sive, H. (2003). *vhnf1* and Fgf signals synergize to specify rhombomere identity in the zebrafish hindbrain. *Development* 130, 3821–3829.
- Wilkinson, D.G., Bhatt, S., Cook, M., Boncinelli, E., and Krumlauf, R. (1989). Segmental expression of Hox-2 homeobox-containing genes in the developing mouse hindbrain. *Nature* 341, 405–409.
- Yashiro, K., Zhao, X., Uehara, M., Yamashita, K., Nishijima, M., Nishino, J., Saijoh, Y., Sakai, Y., and Hamada, H. (2004). Regulation of retinoic acid distribution is required for proximodistal patterning and outgrowth of the developing limb. *Dev. Cell* 6, 411–422.
- Zhang, M., Chen, W.G., Smith, S.M., and Napoli, J.L. (2001). Molecular characterization of a mouse short chain dehydrogenase/reductase active with all-trans-retinol in intact cells, mRDH1. *J. Biol. Chem.* 276, 44083–44090.
- Zhang, M., Hu, P., Krois, C.R., Kane, M.A., and Napoli, J.L. (2007). Altered vitamin A homeostasis and increased size and adiposity in the rdh1-null mouse. *FASEB J.* 21, 2886–2896.